



King's Research Portal

DOI:

[10.1038/eye.2016.149](https://doi.org/10.1038/eye.2016.149)

Document Version

Peer reviewed version

[Link to publication record in King's Research Portal](#)

Citation for published version (APA):

Liew, G., Lee, A. Y., Zarranz-Ventura, J., Stratton, I., Bunce, C., Chakravarthy, U., Lee, C. S., Keane, P. A., Sim, D. A., Akerele, T., McKibbin, M., Downey, L., Natha, S., Bailey, C., Khan, R., Antcliff, R., Armstrong, S., Varma, A., Kumar, V., ... Tufail, A. (2016). The UK Neovascular AMD Database Report 3: inter-centre variation in visual acuity outcomes and establishing real-world measures of care. *Eye (London, England)*, 30(11), 1462-1468. <https://doi.org/10.1038/eye.2016.149>

Citing this paper

Please note that where the full-text provided on King's Research Portal is the Author Accepted Manuscript or Post-Print version this may differ from the final Published version. If citing, it is advised that you check and use the publisher's definitive version for pagination, volume/issue, and date of publication details. And where the final published version is provided on the Research Portal, if citing you are again advised to check the publisher's website for any subsequent corrections.

General rights

Copyright and moral rights for the publications made accessible in the Research Portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognize and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the Research Portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the Research Portal

Take down policy

If you believe that this document breaches copyright please contact librarypure@kcl.ac.uk providing details, and we will remove access to the work immediately and investigate your claim.

The UK Neovascular AMD Database Report 3: Inter-Centre Variation in Visual Acuity Outcomes and Establishing Real-World Measures of Care.

Subtitle: Real world Outcomes in UK centres.

Gerald Liew Phd*, Aaron Y Lee MD*, Javier Zarranz-Ventura PhD, Irene Stratton FFPH, Catey Bunce PhD, Usha Chakravarthy PhD, Cecilia S Lee MD, Pearse A. Keane MD, Dawn Sim PhD, Toks Akerele MD, Martin McKibbin MD, Louise Downey MD, Salim Natha MD, Clare Bailey MD, Rehna Khan MD, Richard Antcliff MD, Stewart Armstrong MD, Atul Varma MD, Vineeth Kumar MD, Marie Tsaloumas MD, Kaveri Mandal MD, Catherine Egan PhD, Robert L Johnston** PhD, Adnan Tufail** PhD, on behalf of UK AMD EMR Users Group

* These authors contributed equally and GL and AYL should be considered joint first authors

** These authors contributed equally and RLJ and AT should be considered joint senior authors

Corresponding author – Adnan Tufail

Moorfields Eye Hospital NHS Trust, 162 City Road, London, EC1V 2PD

Adnan.Tufail@moorfields.nhs.uk

Conflict of interest. Robert L Johnston is the Medical Director of Medisoft Limited, the Electronic Medical Record software provider from which data were extracted. All other authors have no relevant financial conflicts of interest to disclose.

Keywords: ranibizumab; neovascular AMD; Age-related Macular degeneration National Dataset; outcome; audit; benchmark standard; Electronic Medical Record

Word count: 219

- 28 **Complete UK AMD EMR Users Group (listed alphabetically by centre and within each centre)**
- 29 **Belfast Health and Social Care Trust, UK** - Usha Chakravarthy
- 30 **Calderdale and Huddersfield NHS Foundation Trust, UK** - Rehna Khan
- 31 **Countess of Chester Hospital NHS Trust, UK** - Stewart Armstrong
- 32 **Gloucestershire Hospitals NHS Foundation Trust, UK** - Robert Johnston, Quresh Mohamed, Ahmed Sallam,
33 Javier Zarranz-Ventura, Paul Donachie, Irene Stratton
- 34 **Hinchingbrooke health care NHS Trust, UK** - Toks Akerele
- 35 **Hull and East Yorkshire Hospitals NHS Trust, UK** - Seema Arora, Helen Cook, Louise Downey, Kala
36 Gopalakrishnan, Fiona Lyon, Tahir Islam, Naeem Zaman.
- 37 **Leeds Teaching Hospitals NHS Trust, UK** - Oliver Backhouse, Tim Dabbs, Bryn Davies, Martin McKibbin,
38 Bataung Mokete, Damian O'Neill
- 39 **Mid Yorkshire Hospitals NHS Trust, UK** - Atul Varma
- 40 **Moorfields Eye Hospital NHS Trust, UK** - Catey Bunce, Catherine Egan, Pearse A Keane, Gerald Liew,
41 Praveen J Patel, Dawn A Sim, Adnan Tufail, Wen Xing
- 42 **Royal United Hospital Bath NHS Trust, UK** - Richard Antcliff
- 43 **University Hospital Birmingham NHS Foundation Trust, UK** - Marie Tsaloumas
- 44 **University Hospitals Bristol NHS Foundation Trust, UK** - Clare Bailey
- 45 **University of Sydney, Sydney NSW, Australia** – Gerald Liew
- 46 **University of Washington, Washington, USA** – Aaron Lee, Cecilia S Lee
- 47 **Warrington and Halton Hospitals NHS Foundation Trust, UK** - Kaveri Mandal
- 48 **Wirral University Teaching Hospital NHS Foundation Trust, UK** - Vineeth Kumar
- 49 **Wrightington, Wigan and Leigh NHS Foundation Trust, UK** - Salim Natha

50

51 **Financial Disclosures**

52 Robert L Johnston is the Medical Director of Medisoft Limited, the Electronic Medical Record software
53 provider from which data were extracted. All other authors have no relevant financial conflicts of interest
54 to disclose.

55 All other authors state that they have no relevant conflicts of interest to disclose.

56

57

59 **Abstract**

60 **Aims:** International variations in visual acuity (VA) outcomes of eyes treated for neovascular age-related
61 macular degeneration (nAMD) are well documented, but intra-country inter-centre regional variations are
62 not known. This data is important for national quality outcome indicators. We aimed to determine intra-
63 country, inter-centre regional variations in outcomes for treatment of nAMD.

64 **Methods:** Prospective multi-centre national database study of 13 UK centres that treated patients
65 according to a set protocol (3 loading doses, followed by pro-re-nata re-treatment). 5,811 treatment naive
66 eyes of 5,205 patients received a total of 36,206 ranibizumab injections over 12 months.

67 **Results:** Mean starting VA between centres varied from 48.9 to 59.9 ETDRS letters. Mean inter-centre VA
68 change from baseline to 12 months varied from +6.9 letters to -0.6 letters (mean of +2.5 letters). The
69 proportion of eyes achieving VA of 70 letters or more varied between 21.9% and 48.7% at 12 months.
70 Median number of injections (visits) at each centre varied from 5 to 8 (9 to 12) with an overall median of 6
71 (11). Age, starting VA, number of injections and visits but not gender were significantly associated with
72 variation in these VA outcomes ($P<0.01$). Significant variation between centres persisted even after
73 adjusting for these factors.

74 **Conclusions:** There are modest differences in VA outcomes between centres in the UK. These differences
75 are influenced, but not completely explained, by factors such as patient age, starting VA, number of
76 injections and visits. These data provide an indication of the VA outcomes that are achievable in real world
77 settings.

78

79 **Introduction**

80 Age-related macular degeneration (AMD) is one of the leading causes of irreversible blindness in patients
81 aged over 60 years. Since the introduction of anti-Vascular Endothelial Growth Factor (VEGF) agents to
82 treat neovascular AMD (nAMD) in 2006, rates of blindness and visual impairment from AMD have declined
83 dramatically.¹⁻³ Data from several randomized clinical trials suggest patients gain on average 6-11 letters
84 using the most aggressive monthly dosing posology in the first year of treatment.⁴⁻⁷ A less resource
85 intensive Pro-Re-Nata (PRN) dosing posology has been found in the Comparison of Age-related macular
86 degeneration Treatment Trials (CATT)^{6;8} and Inhibit VEGF in Age-related choroidal Neovascularisation
87 (IVAN)^{7;9} trials to produce outcomes similar to monthly dosing. This has provided a sound empirical basis
88 for use of PRN posology in the National Health System (NHS) in the United Kingdom (UK).

89

90 Although PRN dosing posology in theory is able to achieve excellent outcomes, ‘real-world’ studies have not
91 matched the outcomes achieved in clinical trials.¹⁰⁻¹³ We have previously reported longitudinal results from
92 a cohort of patients recruited from hospitals in the UK.^{10;11;14} Patients had standardised data recorded at
93 the point of care into an Electronic Medical Record (EMR) system and the grouped national findings
94 showed that PRN retreatment after 3 loading doses resulted in moderately stable vision.¹¹ National data
95 have also been reported from Australia for a treat and extend posology.¹⁵ International multi-country real
96 life comparisons have reported some international differences in VA outcomes and treatment
97 patterns,¹⁶ which may be influenced by different reimbursement and health system structures. However,
98 what remains unclear is how much, if any, intra-country inter-centre regional variation occurs. This data is
99 obscured by aggregate national data, and is important as it permits understanding of the factors that
100 influence outcomes in a real world setting and allows national quality benchmarks to be set. In this report
101 we examine inter-centre variations in patient characteristics, the number of treatments delivered and their
102 impact on visual outcomes in 13 geographically distinct UK centres.

103

104 **Methods**

105 **Study Design**

106 The study design is described in detail in previous reports from the UK-nAMD Database Study group.^{10;11;14}
107 In brief, sites known to make comprehensive use of Electronic Medical Records (EMR) systems were
108 contacted and requested to contribute data. Patient identifiers were completely removed and site and
109 clinician data were pseudo-anonymised. On this basis an ethics committee determined that formal ethics
110 approval was not required. The study was conducted in accordance with the declaration of Helsinki and the
111 UK's Data Protection Act.

112 **Study Centres**

113 Thirteen NHS hospitals that deliver ranibizumab AMD treatment services in England and Northern Ireland
114 submitted data to this study. Each site is the sole NHS provider of nAMD care to their local population and
115 very few patients switch between providers. Following NICE approval for the use of ranibizumab for nAMD
116 in the NHS in August 2008 all sites used this drug almost exclusively, although prior to this date some sites
117 offered limited treatment with bevacizumab. The study was initiated on 1st Feb 2012, all approvals and data
118 extraction was performed by 02 April 2012. Data was delivered to the analysis team by the end of April
119 2012.

120 **Data variables**

121 Analysis was restricted to treatment-naïve eyes undergoing ranibizumab therapy conforming to 3 monthly
122 loading dose followed by PRN posology for nAMD that completed at least 12 months of follow up.
123 In this report the 'best-measured VA' was the best VA with refraction or habitual correction and/or pinhole
124 as measured on an ETDRS chart and expressed as LogMAR vision and ETDRS letters. The vast majority of
125 sites measured VA with habitual correction rather than best-corrected refracted VA at all time points and
126 used ETDRS charts. Analysis for eyes with very low VA was undertaken by substituting counting fingers (CF),
127 hand movement (HM), and perception of light (PL) with 2.0, 2.3, and 2.7, respectively.
128 We examined the outcome measures commonly reported in clinical trials (VA change from 0 to 12 months,
129 proportion of eyes gaining and losing 15 ETDRS letters etc) as well as other measures such as VA change
130 from 3 to 12 months and proportion of eyes achieving 70 ETDRS letters at 12 months (Snellen equivalent of
131 20/40 or 6/12, driving equivalent in many jurisdictions).

132 **Statistical methods**

133 Medisoft Ophthalmology (Medisoft Limited, Leeds, UK) was the EMR system used for data extraction. Data
134 for right and left eyes of patients who had had at least one intravitreal injection of ranibizumab for nAMD
135 were extracted. Both STATA version 11 and SPSS version 19 were used to analyse data. Perl and R package
136 ggplot2 was used for multivariable analyses, construction of generalised linear models and creation of
137 funnel plots.

138 **Results**

139 **Participants**

140 Over the 1 year of follow up analysed in this study, 36,206 ranibizumab injections were performed in 5,811
141 eyes. Table 1 shows the baseline demographics of each of the 13 centres. The number of eyes treated
142 ranged from 39 (Centre M) to 923 (Centre C). Patients treated at each centre were of similar age, ranging
143 from a mean of 78.3 (J) to 81.7 years at the time of the first injection (K).

144 Treatment and outcome characteristics of each centre are shown in Table 2. The median number of
145 injections in each centre was 6 with one centre providing a median of 8 injections (J) and 2 centres
146 providing a median of 5 (L, M). Mean starting VA across all centres was lowest in centre F (48.9 ETDRS
147 letters) and highest in centre L (59.9 letters). VA after 12 months of treatment was highest in centre I (63.2
148 letters) and lowest in centre M (52.9 letters). The centres with the higher mean starting VA did not
149 necessarily finish 12 months with the highest VA at 1 year, nor did the centres with the lowest starting VA
150 finish with the lowest VA at 1 year. Centres that saw patients the most frequently (B) or injected most
151 frequently (J) achieved the 5th and 2nd best VA at 1 year respectively and showed the least variation
152 between 3 and 12 months.

153 At 12 months, the mean change in VA across 13 centres varied from +6.9 letters (centre I) to -0.6 letters
154 (centre L), a difference of 7.5 letters, with a mean VA gain of +2.5 letters. Figure 1 shows these results
155 graphically, with further details in Table 2. The funnel plot in Figure 1 shows the distribution of centres and
156 95% and 99% confidence intervals. Two centres (L, D) were slightly outside the 99% confidence intervals;
157 these were the only centres reporting a slight reduction in VA. One of these centres had the lowest number
158 of injections (5, L); however the other centre had the median number of injections (6, D). Starting VA in

159 centre D was near the middle of the distribution, at 53.7 letters, while in centre L, eyes commenced
160 treatment with the best starting VA at 59.9 letters. In contrast, Figure 2 (supplement) shows that there was
161 less variation in VA change from 0 to 3 months, with a tighter clustering of results. Figure 3 (supplement)
162 shows the VA change from 3 months to 12 months, with most centres clustered within the 95% confidence
163 intervals, and the previous 2 centres that were outside the 99% CI when considering change from 0-12
164 months are now either within or very close to the limits (L, D).

165 The proportion of eyes that gained 15 ETDRS letters or more at 12 months ranged from 7.7% (A) to 29.5%
166 (K). When examined on the funnel plot (Figure 4 supplement), these proportions showed little variation
167 from centre to centre, with all centres within or above the 99% CI. The proportion of eyes that lost 15
168 ETDRS letters or more at 12 months showed an even tighter distribution, with all centres within the 99% CI,
169 and all except one (L) within the 95% CI (Figure 5 supplement). The actual proportions ranged from 4.6% (I)
170 to 11.7%(L). We also examined the proportion of eyes maintaining or achieving driving vision of 70 ETDRS
171 letters or more and found all centres performed above the lower 99% CI limit.(Figure 6)

172 It should be noted that the centre with the lowest number of visits and injections (M, median 9 visits,
173 median 5 injections, Table 2) had outcomes in the middle of the distributions for all the VA measures
174 studied (Figures 1-6). Similarly, the centres with the highest number of visits (B, 12 visits) and highest
175 number of injections (J, 8 injections), generally had outcomes either in the middle of the distribution or
176 higher than average but still within the 95% CI (Figures 1-6).

177 We performed multivariable analyses to determine which factors were associated with better visual
178 outcomes. Younger age, worse starting VA, and higher number of injections and visits but not gender were
179 significantly associated with variation in these VA outcomes ($P<0.01$). Significant variation between centres
180 persisted even after adjusting for these factors.

181 **Discussion**

182 There is considerable published data on VA outcomes derived from clinical trials but limited data describing
183 real-world outcomes. Real-world outcomes indicating what is possible when trial results translate to clinical
184 practice are ultimately the most important measure as they reflect what happens to whole populations of
185 patients rather than the rarefied cohorts included in trials. They are also important for establishing

186 benchmarks standards that are achievable in busy public systems, and for defining measures of quality care
187 that take into account the heterogeneity of patient populations and care delivery systems. This study
188 provides some of the first real-world outcomes from a single national health system, namely 13 UK public
189 hospital centres using a PRN treatment posology. We report that there was some inter-centre variation in
190 VA outcomes up to a maximum difference of 7.5 letters between the highest and lowest VA achieved from
191 0-12 months. Age, starting VA, number of injections and visits but not gender were significantly associated
192 with variation in these VA outcomes.

193 The median performance of these 13 centres is comparable to results from clinical trials, once the lower
194 starting VA and lower number of injections is taken into account. Table 3 compares findings from this study
195 with clinical trial results. The CATT⁶ and IVAN^{7,9} studies achieved mean improvement from 0-12 months of
196 6.8 and 5.0 letters, respectively with 7 injections, while the Groupe d'Etude Français Avastin versus Lucentis
197 dans la DMLA néovasculaire (GEFAL)¹⁷ and Multicentre Anti-VEGF Trial in Austria (MANTA)¹⁸ studies
198 showed mean improvement of 2.9 and 4.1 letters respectively with 6 injections, as compared with mean
199 improvement of 2.5 letters with 6 injections in this study. These results suggest that a similar benchmark of
200 +2.5 letters improvement (0-12 months) with 6 injections represents quality ongoing care that is achievable
201 in a real-world, public hospital setting.

202 There are few other real-world studies with which to compare our results. A large database observational
203 study from Australia, the Fight Retinal Blindness Study,¹⁹ reported mean VA gains of 5.3 letters after 2 years
204 and 13 injections of a treat and extend posology. These results are superior to those achieved in this report
205 but direct comparisons are difficult due to differences in health systems, patient mix, and different follow-
206 up periods.

207 It should be noted that the difference in 0-12 month VA change between the highest and lowest scoring
208 centres was 7.5 ETDRS letters, a difference that is only marginally beyond what some studies have
209 considered non-inferior. The CATT⁶ and GEFAL¹⁷ considered a difference of 5 EDTRS letters to represent
210 noninferiority, while the MANTA¹⁸ considered a difference of 7 letters to be noninferior.¹⁸

211 Due to the 'ceiling effect' whereby eyes starting treatment with good VA have little room for further
212 improvement, many measures are dependent on the starting VA.^{11;12;20} Adjusting for age, starting VA,
213 number of injections and visits reduced, but did not eliminate the significant variation between centres,
214 suggesting there are other unmeasured factors that contribute to these variations in outcomes.

215

216 This study has several strengths including a large sequential sample, collection of a standardised minimum
217 dataset as mandated by the use of an EMR reflecting routine, real-world clinical practice and the large
218 number of centres involved. A weakness of this study is the loss to follow up of a number of patients over
219 time, as is inevitable in a real-world clinical setting. Although there were differences in baseline
220 demographics between patients lost to follow-up and those who completed follow-up, we have previously
221 shown that VA changes are similar in both groups.¹¹ Best-corrected VAs were not routinely measured,
222 instead the VAs with habitual correction were reported in this study which may underestimate absolute VA
223 measurements compared with clinical trials. It should be noted that clinical treatment decisions were based
224 on these VAs and we believe these represent real-world outcomes and may better reflect patients' visual
225 experience than protocol determined best-corrected VAs. A possible reason for differences in injection
226 numbers may be individual differences in centre / physician thresholds for retreatment using a PRN
227 posology.

228

229 In summary, we report that 13 UK centres using a PRN treatment posology for managing neovascular AMD
230 achieved broadly similar VA outcomes with modest variability in outcomes. The difference between the
231 highest and lowest VA gain at 12 months was 7.5 letters with a mean of +2.5 letters gained. Age, starting
232 VA, number of injections and visits but not gender were significantly associated with variation in these VA
233 outcomes. These data may be considered as establishing an achievable benchmark for the quality of PRN
234 posology in real-world settings, and is likely to be relevant to many sites worldwide.

235

236

237

238 **Acknowledgements.**

239

240 **Funding**

241 This study group has received a proportion of its funding from the Department of Health's NIHR Biomedical
242 Research Centre for Ophthalmology at Moorfields Eye Hospital and UCL Institute of Ophthalmology. Cecelia
243 S Lee is funded by NIH/NEI grant NIH/NEI K23EY024921. The views expressed in the publication are those of
244 the authors and not necessarily those of the Department of Health. The funding sources had no role in the
245 design and conduct of the study; collection, management, analysis, and interpretation of the data;
246 preparation, review, or approval of the manuscript; and decision to submit the manuscript for publication.

247

248 **Financial Disclosures**

249 Robert L Johnston is the Medical Director of Medisoft Limited, the Electronic Medical Record software
250 provider from which data were extracted. All other authors have no relevant financial conflicts of interest
251 to disclose.

252

253 **Other Acknowledgements**

254

255 The authors acknowledge Paul Donachie, Gloucestershire Hospitals NHS Foundation Trust for providing
256 statistical advice on the manuscript.

258

259

260 GL and AF had full access to all the data in the study and take responsibility for the integrity of the data and
261 the accuracy of the data analysis.

262

- 265 1. Bloch SB, Larsen M, Munch IC. Incidence of legal blindness from age-related macular degeneration in
266 denmark: year 2000 to 2010. *Am J Ophthalmol*. 2012; 153:209-213.
- 267 2. Skaat A, Chetrit A, Belkin M, Kinori M, Kalter-Leibovici O. Time trends in the incidence and causes of
268 blindness in Israel. *Am J Ophthalmol*. 2012; 153:214-221.
- 269 3. Buckle M, Donachie PH, Johnston RL. Long-term outcomes of intravitreal ranibizumab for neovascular
270 age-related macular degeneration in a well defined region of the UK. *Br J Ophthalmol*. 2015.
- 271 4. Rosenfeld PJ, Brown DM, Heier JS, Boyer DS, Kaiser PK, Chung CY et al. Ranibizumab for Neovascular
272 Age-Related Macular Degeneration. *N Engl J Med*. 2006; 355:1419-1431.
- 273 5. Brown DM, Kaiser PK, Michels M, Soubrane G, Heier JS, Kim RY et al. Ranibizumab versus Verteporfin
274 for Neovascular Age-Related Macular Degeneration. *N Engl J Med*. 2006; 355:1432-1444.
- 275 6. Martin DF, Maguire MG, Ying GS, Grunwald JE, Fine SL, Jaffe GJ. Ranibizumab and bevacizumab for
276 neovascular age-related macular degeneration. *N Engl J Med*. 2011; 364:1897-1908.
- 277 7. Chakravarthy U, Harding SP, Rogers CA, Downes SM, Lotery AJ, Wordsworth S et al. Ranibizumab
278 versus bevacizumab to treat neovascular age-related macular degeneration: one-year findings from
279 the IVAN randomized trial. *Ophthalmology*. 2012; 119:1399-1411.
- 280 8. Martin DF, Maguire MG, Fine SL, Ying GS, Jaffe GJ, Grunwald JE et al. Ranibizumab and bevacizumab
281 for treatment of neovascular age-related macular degeneration: two-year results. *Ophthalmology*.
282 2012; 119:1388-1398.
- 283 9. Chakravarthy U, Harding SP, Rogers CA, Downes SM, Lotery AJ, Culliford LA et al. Alternative
284 treatments to inhibit VEGF in age-related choroidal neovascularisation: 2-year findings of the IVAN
285 randomised controlled trial. *Lancet*. 2013; 382:1258-1267.
- 286 10. Zarranz-Ventura J, Liew G, Johnston RL, Xing W, Akerele T, McKibbin M et al. The Neovascular Age-
287 Related Macular Degeneration Database: Report 2: Incidence, Management, and Visual Outcomes of
288 Second Treated Eyes. *Ophthalmology*. 2014; 121:1966-1975.
- 289 11. The neovascular age-related macular degeneration database: multicentre study of 92 976
290 ranibizumab injections: report 1: visual acuity. *Ophthalmology*. 2014; 121:1092-1101.
- 291 12. Gabai A, Veritti D, Lanzetta P. One-year outcome of ranibizumab for neovascular age-related macular
292 degeneration: a thorough analysis in a real-world clinical setting. *Eur J Ophthalmol*. 2014; 24:396-401.
- 293 13. Dadgostar H, Ventura AA, Chung JY, Sharma S, Kaiser PK. Evaluation of injection frequency and visual
294 acuity outcomes for ranibizumab monotherapy in exudative age-related macular degeneration.
295 *Ophthalmology*. 2009; 116:1740-1747.
- 296 14. Lee AY, Lee CS, Butt T, Xing W, Johnston RL, Chakravarthy U et al. UK AMD EMR USERS GROUP
297 REPORT V: benefits of initiating ranibizumab therapy for neovascular AMD in eyes with vision better
298 than 6/12. *Br J Ophthalmol*. 2015; 99:1045-1050.
- 299 15. Gillies MC, Walton R, Simpson JM, Arnold JJ, Guymer RH, McAllister IL et al. Prospective audit of
300 exudative age-related macular degeneration: 12-month outcomes in treatment-naïve eyes. *Invest*
301 *Ophthalmol Vis Sci*. 2013; 54:5754-5760.

- 302 16. Holz FG, Tadayoni R, Beatty S, Berger A, Cereda MG, Cortez R et al. Multi-country real-life experience
303 of anti-vascular endothelial growth factor therapy for wet age-related macular degeneration. *Br J*
304 *Ophthalmol*. 2015; 99:220-226.
- 305 17. Kodjikian L, Souied EH, Mimoun G, Mauget-Faysse M, Behar-Cohen F, Decullier E et al. Ranibizumab
306 versus Bevacizumab for Neovascular Age-related Macular Degeneration: Results from the GEFAL
307 Noninferiority Randomized Trial. *Ophthalmology*. 2013; 120:2300-2309.
- 308 18. Krebs I, Schmetterer L, Boltz A, Told R, Vecsei-Marlovits V, Egger S et al. A randomised double-
309 masked trial comparing the visual outcome after treatment with ranibizumab or bevacizumab in
310 patients with neovascular age-related macular degeneration. *Br J Ophthalmol*. 2013; 97:266-271.
- 311 19. Arnold JJ, Campain A, Barthelmes D, Simpson JM, Guymer RH, Hunyor AP et al. Two-year outcomes of
312 "treat and extend" intravitreal therapy for neovascular age-related macular degeneration.
313 *Ophthalmology*. 2015; 122:1212-1219.
- 314 20. Ross AH, Donachie PH, Sallam A, Stratton IM, Mohamed Q, Scanlon PH et al. Which visual acuity
315 measurements define high-quality care for patients with neovascular age-related macular
316 degeneration treated with ranibizumab? *Eye (Lond)*. 2013; 27:56-64.
317
318

319

320 Figure Legends.

321 **Figure 1:** Funnel plot showing the change in ETDRS letters from baseline to 12 months by centre. Solid black
322 lines represent the 95% confidence intervals, dashed lines the 99% confidence intervals. ETDRS refers to
323 Early Treatment Diabetic Retinopathy Study.

324
325 **Figure 2 (supplement):** Funnel plot showing the change in ETDRS letters from baseline to 3 months by
326 centre. Solid black lines represent the 95% confidence intervals, dashed lines the 99% confidence intervals.
327 ETDRS refers to Early Treatment Diabetic Retinopathy Study.

328
329 **Figure 3 (supplement):** Funnel plot showing the change in ETDRS letters from 3 months to 12 months by
330 centre. Solid black lines represent the 95% confidence intervals, dashed lines the 99% confidence intervals.
331 ETDRS refers to Early Treatment Diabetic Retinopathy Study.

332
333 **Figure 4 (supplement):** Funnel plot showing the proportion of eyes gaining 15 ETDRS letters or more from
334 baseline to 12 months by centre. Solid black lines represent the 95% confidence intervals, dashed lines the
335 99% confidence intervals. ETDRS refers to Early Treatment Diabetic Retinopathy Study.

336
337 **Figure 5 (supplement):** Funnel plot showing the proportion of eyes losing 15 ETDRS letters or more from
338 baseline to 12 months by centre. Solid black lines represent the 95% confidence intervals, dashed lines the
339 99% confidence intervals. ETDRS refers to Early Treatment Diabetic Retinopathy Study.

340
341 **Figure 6:** Funnel plot showing the proportion of eyes maintaining driving vision of 70 ETDRS letters or better
342 at 12 months by centre. Solid black lines represent the 95% confidence intervals, dashed lines the 99%
343 confidence intervals. ETDRS refers to Early Treatment Diabetic Retinopathy Study.

344
345
346

347

348

349

350

